

# Oxytocin Compound

Published: March 1, 2026

## What is Oxytocin Compound?

Oxytocin (oks-i-TOE-sin)

- Oxytocin is a hormone used for various indications and conditions determined by your doctor.

**Compounded Drug Forms:** nasal spray, tablet, fast-burst sublingual tablet, suspension, troche, cream, gel,

## What should I tell my health care provider before I take this medicine?

- Pregnant or think you may be pregnant; this medication can cause premature labor.
- Consult with your provider if you are breastfeeding.

## How should I use this medicine?

Follow the package directions provided by Belmar Pharma Solutions and by your prescriber. Your dosage is based on your medical condition and response to therapy. Follow the dosing schedule provided carefully. This medication may be taken on an 'as needed' basis as well.

Nasal sprays may need to be primed. Blow nose gently to clear any mucus from nasal passages and wash hands. Hold the bottle with thumb at the bottom and the nozzle between two fingers (use L hand for R nostril and vice versa). Tilt head forward slightly and close nostril not receiving medication. Insert the tip of the nozzle gently into nostril roughly  $\frac{1}{4}$  to  $\frac{1}{2}$  inch inward. Point spray toward ear, away from nasal septum at center. Sniff deeply while pushing down on nozzle once. Breathe out through the mouth. Repeat with other nostril if dosing suggests waiting 10 seconds between sprays.

Oral dosage forms may be taken with or without food. If it upsets your stomach take it with a small meal. Pick a convenient time of day to take your dose and try to take your medication as close to that time each day as possible.

Sublingual formulations should be placed under the tongue or between the cheek and gums and held in place until fully dissolved. Avoid swallowing saliva to ensure best absorption into the bloodstream. Avoid eating or drinking 15 minutes before or after taking sublingual formulations.

Topical products can be applied to the inner arm, upper thigh, back of the knee, tops of the feet and

inner wrists. It is recommended to rotate sites to ensure even retention of the hormone. Cover treated skin with clothes to avoid contact with people or pets.

Vaginal preparations should be inserted using your finger or the applicator provided. Suppositories may leak, and may be more convenient to use at bedtime.

## **What if I miss a dose?**

- If you miss a dose, take it as soon as possible.
- If it is almost time for your next dose, skip the missed dose and continue with your regular dosing schedule.
- Do not take double the dose at once.

## **What should I watch for while using this medicine?**

Belmar Pharma Solutions recommends yearly checkups with your provider.

## **What side effects may I notice from receiving this medicine?**

Possible side effects that may occur and may go away as your body gets used to the medication include:

- Nausea or vomiting
- Headache
- Tingling/itching of the nipples
- Breast tenderness
- Flushing

### **Serious Side Effects:**

Contact your doctor immediately or seek medical attention if you experience:

- Change in heart rate
  - Risk of cardiovascular effects is dependent on dose and route of administration and may be higher in patients with cardiovascular disease.
- Signs of high blood pressure (bad headache, dizziness, passing out, change in eyesight)
- Unusual swelling or weight gain that may be associated with water retention

- Trouble passing urine
- Severe stomach pain
- Seizures
- Weakness on one side of the body
- Trouble speaking or thinking
- Excessive bleeding (soaking 1 pad per hour)

You may experience side effects not listed. Contact your doctor with questions.

An allergic reaction to this medication is unlikely but seek medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, trouble breathing.

## Where should I keep my medicine?

- Follow the storage recommendations found on your prescription label.
- Store at room temperature between 59- and 86-degrees F, unless otherwise specified on the packaging.

### **Additional Information:**

- Do not share this medication with others.
- Keep out of reach of children and pets.
- If an overdose is suspected contact your local poison control center at 1-800-222-1222 or go to the nearest emergency room.

If you have any questions concerning this or any of your medications, a Pharmacist is available during normal business hours to answer your questions. You may report side effects to the FDA at 1-800-FDA-1088.

### **Consumer Information Use and Disclaimer:**

This information should not be used to decide whether or not to take this medication or any other medication. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does not include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. For questions about the source of any of the information provided herein, pharmacists at

Belmar Pharma Solutions are available. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of this medicine.

**NIOSH Drug: Observe special handling and disposal requirements.**

**Do not flush down a toilet or pour down a drain. Check with your pharmacist if you have questions about the best way to dispose of medications. There may be a drug take-back program in your area.**

**References:**

1. Oxytocin. Lexi-Drugs. Lexicomp. Wolters Kluwer Health, Inc. Riverwoods, IL. Available at: <http://online.lexi.com>. Accessed March 2023.

[Learn More About This Medication](#)